

GROUP GRADINGS ON FINITE DIMENSIONAL LIE ALGEBRAS

DUŠAN PAGON, DUŠAN REPOVŠ, AND MIKHAIL ZAICEV

ABSTRACT. We study gradings by noncommutative groups on finite dimensional Lie algebras over an algebraically closed field of characteristic zero. It is shown that if L is graded by a non-abelian finite group G then the solvable radical R of L is G -graded and there exists a Levi subalgebra $B = H_1 \oplus \cdots \oplus H_m$ homogeneous in G -grading with graded simple summands $H_1, \dots, H_m$. All supports $\text{Supp } H_i, i = 1 \dots, m$, are commutative subsets of G .

1. INTRODUCTION

Graded rings and graded algebras have been extensively studied during the last decades (cf. e.g. [2] - [11]). Group gradings were investigated both in the associative case ([5] - [8], [16]) and in the Lie case ([3],[11],[12],[15],[17]), or in other nonassociative cases ([2],[3],[9]).

One of the important tasks is the description of all possible gradings on different algebras. For example, one of the well-known and actively used results in Lie theory is the description of $\mathbb{Z}$ -gradings on finite-dimensional complex Lie algebras [12]. The description of all $\mathbb{Z}_2$ -gradings on matrix algebras plays an exceptional role in the theory of algebras with polynomial identities (cf. [10]).

All abelian gradings on matrix algebras were described in [6]. It was found that the so-called fine gradings on full matrix algebra play an exceptional role in the theory of orthogonal Cartan decompositions of simple complex Lie algebras (cf. [13]). For the noncommutative case all group gradings on matrices were described in [5]. In particular, it was shown that in general case fine gradings are closely connected with some problems of the theory of projective representations of finite groups.

All finite dimensional graded simple associative algebras were described in [7]. This description is presently being very intensively applied in PI-theory (cf. e.g. [1], [14]). All finite dimensional $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded simple Lie algebras were described in [2] and this result was applied for the classification of simple color Lie superalgebras. So, the description of all possible gradings on algebras plays an important role both in the structure theory of finite dimensional and infinite dimensional graded algebras and in its applications.

In the present paper we shall study general properties of commutative and noncommutative gradings on finite dimensional Lie algebras over an algebraically

2010 *Mathematics Subject Classification.* Primary 17B70; Secondary 17B05.

Key words and phrases. Lie algebra, group grading, noncommutative group.

This research was supported by the Slovenian Research Agency grants P1-0292-0101, J1-2057-0101, and J1-9643-0101. The third author was partially supported by RFBR grants No 09-01-00303, 09-0190416-Ukr-f-a and SSC-1983.2008.1.

closed field of characteristic zero. First, we shall show that the description of non-commutative gradings on semisimple algebras can be reduced to abelian gradings and to the classification of graded simple algebras (cf. Propositions 2.3 and 3.1.) Then we shall characterize finite dimensional graded simple algebras and show that all of them can be split into four classical series A, B, C, D and five exceptional series, according to the classification of simple Lie algebras (cf. Proposition 3.1 and Remark 3.2). Finally, we shall show that in the case of finite groups, any graded Lie algebra is a split extension of a homogeneous Levi subalgebra and a solvable radical (cf. Theorem 3.4).

2. GENERAL PROPERTIES OF NONCOMMUTATIVE GRADINGS

Let G be a group. Given a Lie algebra L over a field F , we say that L is G -graded if it can be decomposed into a direct sum of subspaces

$$L = \bigoplus_{g \in G} L_g$$

such that $[L_g, L_h] \subseteq L_{gh}$, for any $g, h \in G$. Subspaces L_g are called homogeneous components and an element $x \in L$ is called homogeneous if $x \in L_g$, for some $g \in G$. In this case we write for shortness $\deg x = g$. A subspace (resp. subalgebra, ideal) V is said to be a graded subspace (resp. subalgebra, ideal) if

$$V = \bigoplus_{g \in G} (V \cap L_g).$$

In other words, if $x = x_{g_1} + \dots + x_{g_n}$, where $\deg x_{g_i} = g_i$ and $g_1, \dots, g_n \in G$ are pairwise distinct, then $x \in V$ if and only if all $x_{g_1}, \dots, x_{g_n}$ belong to V .

A subset

$$\text{Supp } L = \{g \in G \mid L_g \neq 0\}$$

is called the support of the grading. Actually, we can suppose that $\text{Supp } L$ generates G . We call an algebra L graded simple if $[L, L] \neq 0$ and L does not contain any non-trivial graded ideals.

We shall usually use the notation $[x_1, \dots, x_n]$ for the left-normed product in a Lie algebra. That is, $[x_1, \dots, x_n] = [[x_1, \dots, x_{n-1}], x_n]$ for all $n \geq 3$. Similarly, we shall write $[W_1, \dots, W_n] = [W_1, \dots, W_{n-1}, W_n]$ for any subspaces $W_1, \dots, W_n \subseteq L$.

It is well-known that in the case of a simple Lie algebra $L = \bigoplus_{g \in G} L_g$, the group G must be commutative (cf. e.g. [15]). First, we generalize this property to the case of graded simple Lie algebras. In the following two lemmas and Proposition 2.3, G is an arbitrary group and L is not necessarily a finite dimensional Lie algebra over an arbitrary field F .

Lemma 2.1. *Let $L = \bigoplus_{g \in G} L_g$ be a G -graded Lie algebra and let $[L_{g_1}, \dots, L_{g_m}] \neq 0$, for some $g_1, \dots, g_m \in G$. Then $g_1, \dots, g_m$ commute in G .*

Proof. First, note that the inequality $[L_g, L_h] \neq 0$ implies $gh = hg$ since $[L_g, L_h] = [L_h, L_g] \subseteq L_{gh} \cap L_{hg}$. Hence our statement is obvious for $m = 2$. Now we apply the induction on m . Suppose that $[x_1, \dots, x_m] \neq 0$, for some $x_1 \in L_{g_1}, \dots, x_m \in L_{g_m}$ and $m \geq 3$. Since $[x_1, \dots, x_m] = [u, x_{m-1}] + [v, [x_{m-1}, x_m]]$, where $u = [x_1, \dots, x_{m-2}, x_m]$ and $v = [x_1, \dots, x_{m-2}]$ (or $v = x_1$ in case $m = 3$) either $[u, x_{m-1}] \neq 0$ or $[v, [x_{m-1}, x_m]] \neq 0$.

In the first case $g_1, \dots, g_{m-2}, g_m$ commute by the inductive hypothesis and g_{m-1} commutes with the product $g_1 \cdots g_{m-2} g_m$. Since $g_1, \dots, g_{m-1}$ also commute, by induction we have $g_m g_{m-1} = g_{m-1} g_m$ that is, g_m commutes with all $g_1, \dots, g_{m-1}$.

In the second case, $g_1, \dots, g_{m-1}$ commute and the product $g_{m-1} g_m$ commutes with all $g_i, 1 \leq i \leq m-2$. Since $g_i g_{m-1} = g_{m-1} g_i, 1 \leq i \leq m-2$, we obtain $g_i g_m = g_m g_i$, for all $i = 1, \dots, m-2$. Clearly, $g_m g_{m-1} = g_{m-1} g_m$ and we have thus completed the proof. $\square$

Lemma 2.2. *Let $L = \oplus_{g \in G} L_g$ be a G -graded Lie algebra and let $gh \neq hg$ for some $g, h \in \text{Supp } L$. Then $I = [Id(L_g), Id(L_h)] = 0$, where $Id(L_g)$ is the ideal of L generated by L_g .*

Proof. Any element of I can be written as a linear combination of products

$$[x, y_1, \dots, y_k, [z, t_1, \dots, t_m]], \quad k \geq 0, m \geq 0,$$

where $x \in L_g, z \in L_h$ and $y_1, \dots, y_k, t_1, \dots, t_m$ are homogeneous elements from L . Now the statement easily follows by Lemma 2.1. $\square$

As an immediate consequence of these remarks we get the following:

Proposition 2.3. *Let $L = \oplus_{g \in G} L_g$ be a G -graded simple Lie algebra. Then $\text{Supp } L$ generates an abelian subgroup of G .*

$\square$

In particular, if L is simple in the non-graded sense then $\text{Supp } L$ is a commutative subset of G . Note that in general, this property does not hold for semisimple algebras. For example, let $L = B_1 \oplus B_2$ be the direct sum of two simple algebras isomorphic to $sl_2(F)$. Given a group G of order 2, $G = \{e, g\}$, we can define G -grading on $H = sl_2(F)$ by setting

$$L_e = \text{Span} \left\{ \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \right\}, \quad L_g = \text{Span} \left\{ \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \right\}.$$

It is now sufficient to take any group G with $gh \neq hg, g, h \in G, g^2 = h^2 = e$, and define G -grading on B_i , using $H_i, i = 1, 2$. Later we shall show that all noncommutative gradings on semisimple Lie algebras are of similar type.

3. STRUCTURE OF FINITE DIMENSIONAL GRADED ALGEBRAS

Now let G be a finite abelian group and let F be an algebraically closed field of characteristic zero. Recall a well-known duality between gradings and automorphism actions on L (cf. e.g. [6]). Let $\widehat{G}$ be the dual group for G , that is the group of all irreducible characters on G . Since G is finite abelian, the group $\widehat{G}$ is isomorphic to G . If $L = \oplus_{g \in G} L_g$ is a G -graded algebra then any $\chi \in \widehat{G}$ acts on L by the automorphism

$$\chi * x_g = \chi(g) x_g,$$

where $x_g \in L_g$ is a homogeneous element of degree g .

Clearly, all subspaces L_g are stable under the $\widehat{G}$ -action. Moreover, a subspace $V \subseteq L$ is graded if and only if V is $\widehat{G}$ -stable. Conversely, if one defines the $\widehat{G}$ -action on L by automorphisms then L can be decomposed into a direct sum

$$(1) \quad L = \bigoplus_{g \in G} L_g$$

where

$$L_g = \{v \in L \mid \chi * v = \chi(g)v \text{ for all } \chi \in \widehat{G}\}$$

and the decomposition (1) is a G -grading on L .

If G is an infinite cyclic group generated by t then we can also define an action of the infinite cyclic group generated by χ on $L = \bigoplus_{g \in G} L_g$, by setting

$$\chi^n * v = \lambda^{nk} v$$

as soon as $v \in L_{t^k}$ where $\lambda \in F^*$ is a fixed element of infinite order. As before, a subspace V of L is graded if and only if $\chi * V = V$. If L is finite dimensional and $\text{Supp } L$ generates G then G is a finitely generated abelian group and we can again identify G -grading of L with the G -action on L .

Using this duality we get the following result.

Proposition 3.1. *Let $L = \bigoplus_{g \in G} L_g$ be a finite dimensional G -graded Lie algebra over an algebraically closed field of characteristic zero.*

- 1) *If L is graded simple then G is abelian and $L = B_1 \oplus \cdots \oplus B_n$ is semisimple with isomorphic simple components $B_1, \dots, B_n$.*
- 2) *If L is semisimple then $L = A_1 \oplus \cdots \oplus A_m$ is a direct sum of graded simple components and $\text{Supp } A_i$ is a commutative subset of G , for any $1 \leq i \leq m$.*

Proof. First let L be graded simple. Then G is abelian by Proposition 2.3. Suppose that L is not semisimple. Then due to the duality between G -gradings and G -action its solvable radical R is a graded ideal since R is stable under the action of any automorphism of L . Hence $R = L$ and $L^2 = [R, R] \neq L$ is also a graded ideal and $L^2 \neq 0$ by the definition of graded simplicity, a contradiction. Hence L is semisimple. Consider the decomposition $L = B_1 \oplus \cdots \oplus B_n$ into a direct sum of simple ideals.

Let $f : L \rightarrow L$ be an automorphism of L . Clearly, $f(B_1) = B_j$ for some $1 \leq j \leq n$. Due to the duality between G -grading and G -action, the orbit of B_1 under the G -action contains all summands $B_1, \dots, B_n$ since L is graded simple. In particular, all $B_1, \dots, B_n$ are isomorphic, and we have proved assertion (1) of the proposition.

Now let L be a semisimple G -graded algebra. As before, L is a direct sum of minimal ideals, $L = B_1 \oplus \cdots \oplus B_n$. Consider the minimal graded ideal A_1 of L . If $A_1 = L$ then there is nothing to prove. If $A_1 \neq L$ then A_1 is a sum of some B_i , say, $A_1 = B_1 \oplus \cdots \oplus B_t$. We shall prove that $B_{t+1} \oplus \cdots \oplus B_n$ is also a graded ideal of L .

First note that the centralizer of any homogeneous element in L is a graded subalgebra of L . Indeed, let

$$(2) \quad [a_h, b_{g_1} + \cdots + b_{g_m}] = 0$$

for some $a_h \in L_h, b_{g_1} \in L_{g_1}, \dots, b_{g_m} \in L_{g_m}$ with pairwise distinct $g_1, \dots, g_m \in G$. If $[a_h, b_{g_i}] = 0$, for all $i = 1, \dots, m$ then we are done. Otherwise fix all $g_{i_1}, \dots, g_{i_k}$ such that $c_1 = [a_h, b_{g_{i_1}}] \neq 0, \dots, c_k = [a_h, b_{g_{i_k}}] \neq 0$. Then h commutes with $g_{i_1}, \dots, g_{i_k}$ by Lemma 2.1, $c_1 \in L_{hg_{i_1}}, \dots, c_k \in L_{hg_{i_k}}$ and all $hg_{i_1}, \dots, hg_{i_k}$ are distinct. On the other hand, $c_1 + \cdots + c_k = 0$, a contradiction. Hence the equality (2) implies that all $b_{g_1}, \dots, b_{g_m}$ commute with a_h .

Now we observe that the intersection of two graded subalgebras is also a graded subalgebra. In particular, the centralizer of A_1 is a graded subalgebra in L . However,

the centralizer of A_1 is $B_{t+1} \oplus \cdots \oplus B_n$ and we complete the proof of the proposition by induction on dimension of L . $\square$

Remark 3.2. *It follows by the previous proposition that any finite dimensional graded simple algebra is associated with one of the finite dimensional simple Lie algebras. In particular, we can say that the graded simple algebra is an algebra one of the types A_l ($l \geq 1$), B_l ($l \geq 2$), C_l ($l \geq 3$), D_l ($l \geq 4$), G_2 , F_4 , E_6 , E_7 , E_8 .*

Now we clarify the structure of a finite dimensional Lie algebra graded by a finite group.

Proposition 3.3. *Let G be a finite group and let $L = \bigoplus_{g \in G} L_g$ be a finite dimensional G -graded Lie algebra over an algebraically closed field of characteristic zero. Then the radical $R = \text{Rad } L$ of L is a graded ideal and there exists a split extension $L = B + R$, where B is a maximal semisimple subalgebra of L homogeneous in G -grading.*

Proof. If G is an abelian group then, as it was mentioned in the proof of Proposition 3.1, the solvable radical of L is graded.

Now let G be a noncommutative group. Consider a decomposition $L = B + R$, where B is a maximal semisimple subalgebra of L , $B = B_1 \oplus \cdots \oplus B_n$ and $B_1, \dots, B_n$ are minimal ideals of B . By Proposition 2.3, algebra L cannot be graded simple. Consider a maximal proper graded ideal P of L .

First suppose that $B \subseteq P$. Then $\dim L/P = 1$. Denote $R_0 = \text{Rad } P$. Then $R_0 \subset R$ and $\dim R/R_0 = 1$. Moreover, $R^2 \subseteq R_0$ and $[B, R_0] \subseteq R_0$. In particular, R_0 is an ideal of L . Since $\dim P < \dim L$, we may suppose by induction that R_0 is a graded ideal of P and L . If $R_0 \neq 0$ then also by induction, R/R_0 is a graded ideal of L/R_0 , hence R is a graded ideal of L .

In case $R_0 = 0$ we have $\dim R = 1$ and hence it is a trivial B -module. Since $[R, R] = 0$ we conclude that R is the center of L . By Lemma 2.1, the center of a graded algebra is homogeneous and we have completed the proof that R is a graded ideal in this case.

Now let $B \not\subseteq P$. Then L/P is graded simple and by Proposition 3.1, it is semisimple, i.e. $R \subseteq P$. Applying the induction on dimension we again conclude that R is a homogeneous subspace of P and hence of L .

Now we prove the existence of Levi subalgebra homogeneous in G -grading. In case G is abelian we apply the duality between G -grading and G -action by automorphisms on L . By [18] there exists a maximal semisimple subalgebra B stable under $\widehat{G}$ -action. So, B is a graded Levi subalgebra of L .

In the general case we consider graded factor-algebra L/R . By Proposition 3.1 we have

$$L/R = \bar{A}_1 \oplus \cdots \oplus \bar{A}_m$$

where any $\bar{A}_i$ is graded simple and $S_i = \text{Supp } \bar{A}_i$ is a commutative subset of G .

Denote by A_i the full preimage of $\bar{A}_i$ over R . Then A_i is graded and $\text{Supp } A_i \supset S_i$. Now we take a subalgebra C_i of A_i generated by all homogeneous $x \in A_i$ with $\deg x \in S_i$. Since S_i is commutative, by the previous remark C_i contains a homogeneous semisimple subalgebra B_i isomorphic $\bar{A}_i$. Note that $\dim(B_1 + \cdots + B_m) = \dim L/R$ hence $B = B_1 \oplus \cdots \oplus B_m$ is a homogeneous Levi subalgebra and we have completed the proof. $\square$

Combining Propositions 2.3, 3.1 and 3.3 we immediately obtain the following:

Theorem 3.4. *Let $L = \bigoplus_{g \in G} L_g$ be a finite dimensional Lie algebra over an algebraically closed field of characteristic zero graded by a finite group G . Then its solvable radical is homogeneous in G -grading and there exists a Levi subalgebra B homogeneous in G -grading. Moreover, B is a direct sum $B = H_1 \oplus \cdots \oplus H_m$ where any H_i is graded simple subalgebra with commutative support $\text{Supp } H_i$ and in the non-graded sense H_i is a direct sum of isomorphic simple components.*

□

REFERENCES

- [1] E. Aljadeff, D. Haile, and M. Natapov, *Graded identities of matrix algebras and the universal graded algebra*, Trans. Amer. Math. Soc. **362** (2010), 3125–3147.
- [2] Yu. Bahturin and D. Pagon, *Classifying color Lie superalgebras*, Contemp. Math. **483** (2009), 37–54.
- [3] Yu. Bahturin and I. Shestakov, *Gradings of simple Jordan algebras and their relation to the gradings of simple associative algebras*, Comm. Algebra **999** (2002), 111–222.
- [4] Yu. A. Bahturin, I.P. Shestakov, and M.V. Zaicev, *Gradings on simple Jordan and Lie algebras*, J. Algebra **283** (2005), 849–868.
- [5] Yu. Bahturin and M. Zaicev, *Group gradings on matrix algebras*, Canad. Math. Bull. **45** (2002), 499–508.
- [6] Yu. A. Bahturin, M. V. Zaicev, and S. K. Sehgal, *Group gradings on associative algebras*, J. Algebra **241** (2001), 677–698.
- [7] Yu. A. Bahturin, M.V. Zaicev, and S.K. Sehgal, *Finite dimensional graded simple algebras*, Matem. Sbornik **199** (2008), 21–40 (Russian).
- [8] M. Cohen and S. Montgomery, *Group-graded rings, smash products, and group actions*, Trans. Amer. Math. Soc. **282**:1 (1984), 237–258.
- [9] A. Elduque, *Gradings on octonions*, J. Algebra **207**(1998), 342–354.
- [10] A. Giambruno and M. Zaicev, *Polynomial identities and asymptotic methods*, Math. Surveys and Monographs vol. **122**, Amer. Math. Soc., Providence, RI, 2005.
- [11] V. G. Kac, *Graded Lie algebras and symmetric spaces*, Funkcional. Anal. Prilozhen. **2** (1968), 93–94.
- [12] I. L. Kantor, *Some generalizations of Jordan algebras*, Trudy Sem. Vekt. Tenz. Anal. **16**(1972), 407–499.
- [13] A. I. Kostrikin and Pham Huu Tiep, *Classification of irreducible decompositions of simple Lie algebras of type A_n* , Algebra i Analiz **3**:3 (1991), 806–109; English transl. in: St. Petersburg Math. J. **3**:3 (1992), 571–593.
- [14] P. Koshlukov and M. Zaicev, *Identities and isomorphisms of graded simple algebras*, Lin. Alg. Appl. **432**:12 (2010), 3141–3148.
- [15] J. Patera and H. Zassenhaus, *On Lie gradings. I*, Linear Algebra Appl. **112** (1989), 87–159.
- [16] O. N. Smirnov, *Simple associative algebras with finite $\mathbb{Z}$ -grading*, J. Algebra **196** (1997), 171–184.
- [17] O. N. Smirnov, *Finite $\mathbb{Z}$ -grading on Lie algebras and symplectic involution*, J. Algebra **218** (1999), 246–275.
- [18] E. J. Taft, *Invariant Wedderburn factors*, Illinois J. Math. **1** (1957), 565–573.

DUŠAN PAGON, FACULTY OF NATURAL SCIENCES AND MATHEMATICS, UNIVERSITY OF MARIBOR, GOSPOSVETSKA 84, MARIBOR, 2000, SLOVENIA

E-mail address: dusan.pagon@uni-mb.si

DUŠAN REPOVŠ, FACULTY OF MATHEMATICS AND PHYSICS, AND FACULTY OF EDUCATION, UNIVERSITY OF LJUBLJANA, P. O. B. 2964, LJUBLJANA, 1001, SLOVENIA

E-mail address: dusan.repovs@guest.arnes.si

MIKHAIL ZAICEV, DEPARTMENT OF ALGEBRA, FACULTY OF MATHEMATICS AND MECHANICS, MOSCOW STATE UNIVERSITY, MOSCOW, 119992 RUSSIA, ZAICEVMV@MAIL.RU